

RX J0123.4-7321, a Be/X-ray binary in the wing of the SMC

R. Sturm¹, F. Haberl¹, W. Pietsch¹, and A. Udalski²

¹ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany

² Warsaw University Observatory, Aleje Ujazdowskie 4, 00-478 Warsaw, Poland

Received 8 November 2012 / Accepted 13 January 2013

ABSTRACT

Aims. To confirm faint Be/X-ray binary candidates from the XMM-*Newton* survey of the Small Magellanic Cloud, we searched for X-ray outbursts in archival ROSAT observations. We found that RX J0123.4-7321 was much brighter when detected with ROSAT than seen 16 years later by XMM-*Newton*.

Methods. We analysed the ROSAT observations and the OGLE *I*-band light curve of the optical counterpart to investigate the nature of the system.

Results. High long-term variability in the X-ray flux of a factor of ~ 150 was found between the ROSAT and XMM-*Newton* detections, indicating strong outburst activity during the ROSAT observations. The *I*-band light curve reveals long-term variability and regular outbursts with a period of (119.9 ± 2.5) days indicating the orbital period of the binary system.

Conclusions. The large X-ray flux variations and the properties of the optical counterpart confirm RX J0123.4-7321 as a new Be/X-ray binary in the wing of the Small Magellanic Cloud.

Key words. galaxies: individual: Small Magellanic Cloud – galaxies: stellar content – stars: emission-line, Be – stars: neutron – X-rays: binaries

1. Introduction

Be/X-ray binaries (BeXRBs, e.g. Reig 2011) are binary systems composed of a compact object and a Be star of spectral type O9–B5 and luminosity class III–V. Be stars eject matter in the equatorial plane leading to the build up of a decretion disc (Okazaki 2001; Ziolkowski 2002) around the star that produces emission lines and a near-infrared (NIR) excess. Both components show variability due to disc instabilities and sometimes due to interaction with the compact object. The compact object is in most cases a neutron star (NS). Systems with black holes are not known up to now. However, systems may contain white dwarfs (Li et al. 2012; Sturm et al. 2012a) that are observable at SMC distance as super-soft X-ray source in the case of thermonuclear surface burning on the white dwarf. Caused by a kick experienced by the NS during the supernova explosion, the NS can have an eccentric orbit around the Be star. Accretion is possible, when the NS is close to the decretion disc of the Be star during periastron passage. This causes so-called type-I outbursts that last for some days. Longer (several weeks) type-II outbursts are likely connected with decretion disc instabilities (Ziolkowski 2002).

The Small Magellanic Cloud (SMC) hosts a large population of BeXRBs. A literature search (e.g. Haberl & Pietsch 2004; Coe et al. 2005; Liu et al. 2005; Laycock et al. 2010) reveals about 60 pulsars in more than 100 BeXRB systems (including candidates), which allows a comparison of their statistical properties with those of the Galactic sample. Most of the BeXRBs in the SMC are found in the bar where increased star formation ~ 40 Myr ago created them in large number (Antoniou et al. 2010). However, in the eastern wing of the SMC only a few systems are known, probably due to a younger stellar population (~ 10 Myr, Harris & Zaritsky 2004). Also the BeXRBs in the wing might be younger, as e.g. SXP 1062 found in a supernova rem-

nant suggests (Hénault-Brunet et al. 2012; Haberl et al. 2012b). This might cause different properties of the wing sample. E.g. a literature search reveals in the wing area three known BeXRB pulsars with $P_{\text{spin}} < 40$ s (Clark et al. 1997; Chakrabarty et al. 1998; Corbet et al. 2003) compared to only one (SXP 1062) pulsar with $P_{\text{spin}} > 40$ s. In contrast to that, for the whole SMC Knigge et al. (2011) find a bimodal spin distribution with more systems above $P_{\text{spin}} > 40$ s, which is likely caused by a bimodal distribution of orbital periods. The search for additional systems in the wing of the SMC is essential to enable a statistical comparison of the wing and bar population and exclude any observational bias.

The XMM-*Newton* survey of the SMC (Haberl et al. 2012a) revealed new BeXRBs in X-ray-bright state (Townsend et al. 2011; Sturm et al. 2011; Coe et al. 2011, 2012). In addition, several candidates for BeXRBs were found in the XMM-*Newton* SMC point-source catalogue (Sturm et al. 2013). One of these candidates is RX J0123.4-7321 (Haberl et al. 2000), which was detected with ROSAT (Trümper 1982) previously at much higher X-ray luminosity, but not recognised as BeXRB. In this paper, we present our analysis of the ROSAT observations and the OGLE (Udalski et al. 2008) data of the optical counterpart, allowing us to confirm RX J0123.4-7321 as a new BeXRB in the wing of the SMC with a determined orbital period.

2. Analyses and results of X-ray observations

RX J0123.4-7321 was detected four times in X-ray observations as listed in Table 1. The source was found in three ROSAT observations of SMC X-1 performed between October 1991 and June 1993 at an off-axis angle of $\sim 27.7'$. The ROSAT position from Haberl et al. (2000, their source 460) is RA (J2000)= $01^{\text{h}}23^{\text{m}}27^{\text{s}}.2$ and Dec (J2000)= $-73^{\circ}21'25''$ with a 90% uncertainty of $4.6''$. This is consistent with the XMM-

Table 1. X-ray observations of RX J0123.4-7321.

Satellite Instrument	ObsID	MJD ^a	Dur ^b (d)	Orbit ^c Phase	Exp (ks)	Cts	F_{observed} ($\text{erg cm}^{-2} \text{s}^{-1}$) 0.1–2.4 keV	F_{model} ($\text{erg cm}^{-2} \text{s}^{-1}$) 0.2–4.5 keV	L^d (erg s^{-1}) 0.1–2.4 keV	L^d (erg s^{-1}) 0.2–4.5 keV
ROSAT/PSPC	rp400022n00	48536.2	0.96	0.65	16.6	936	8.84×10^{-13}	1.96×10^{-12}	6.43×10^{35}	1.07×10^{36}
ROSAT/PSPC	rp400022a01	48895.7	1.87	0.66	9.0	584	1.02×10^{-12}	2.27×10^{-12}	7.43×10^{35}	1.24×10^{36}
ROSAT/PSPC	rp400022a02	49141.0	1.20	0.70	11.9	147	1.95×10^{-13}	4.32×10^{-13}	1.42×10^{35}	2.36×10^{35}
XMM/EPIC	0601212401	55011.6	0.41	0.71	25.4	58	–	1.55×10^{-14}	–	7.86×10^{33}

Notes. ^(a) Modified Julian date of the beginning of the observation. ^(b) Duration of the observation. ^(c) Phase = 0.5 corresponds to maximum of optical outbursts. ^(d) Unabsorbed luminosity. A source distance of 60 kpc is assumed.




Fig. 1. Merged ROSAT/PSPC spectrum of RX J0123.4-7321 from observations rp400022n00 and rp400022a01 together with the best-fit power-law model. The lower panel gives the residuals in units of σ .

Newton position of the SMC-survey catalogue (Sturm et al. 2013, source 3003) of RA (J2000)=01^h23^m27^s.46 and Dec (J2000)=−73°21′23″.4 with a 1σ uncertainty of $1.1''$.

ROSAT spectra were extracted using the *ftools* (Blackburn 1995) task *xselect*. For spectral analysis, we merged the observations rp400022n00 and rp400022a01 to increase the statistics and used *xspec* (Arnaud 1996) version 12.7.0u to fit the spectrum with a power-law model and photoelectric absorption by a Galactic column density set to $N_{\text{H,gal}} = 3.3 \times 10^{20} \text{ cm}^{-2}$ (a foreground H I map was kindly provided by Erik Muller, see also Muller et al. 2003) and an additional column density, accounting for the interstellar medium of the SMC and source intrinsic absorption. We find a good description of the data by this model with $\chi^2/\text{dof}=14.8/17$. For the SMC and source intrinsic absorption we obtain $N_{\text{H,smc}} = 4.32^{+0.36}_{-0.28} \times 10^{21} \text{ cm}^{-2}$. The total line-of-sight $N_{\text{H,smc}}$ column density at the position of RX J0123.4-7321 is $3.02 \times 10^{21} \text{ cm}^{-2}$ (Stanimirovic et al. 1999), pointing to some intrinsic absorption in the binary system. The photon index is determined to be $\Gamma = 1.18^{+0.63}_{-0.54}$, which suggests a neutron-star BeXRB ($\Gamma \approx 1$, Haberl & Pietsch 2004) but we cannot exclude an AGN ($\Gamma \approx 1.75$, Tozzi et al. 2006). The ROSAT spectrum together with the best-fit model is presented in Fig. 1.

The XMM-*Newton* detection is based on only 58 counts. However, hardness ratios are consistent with a BeXRB nature of the source. Assuming the same spectral shape for the ROSAT and XMM-*Newton* detections and comparing their fluxes, we




Fig. 2. OGLE III *I*-band finding chart of RX J0123.4-7321, marked with a white cross. Circles indicate the 90% position uncertainties of the ROSAT and XMM-*Newton* detections.

find a high long-term variability of a factor of 146 ± 20 , that would be atypical for an AGN.

We used a Bayesian odds ratio (Gregory & Loredo 1996; Haberl et al. 2008) and a Rayleigh Z_1^2 (Buccheri et al. 1983; Haberl & Zavlin 2002) test to search for the spin period of the NS. The search for pulsations in the ROSAT data with best statistics (rp400022n00) in the (1–200) s band did not reveal convincing periodicities. Note that the ROSAT observation was performed with individual exposures between 200 and 1300 s, preventing the detection of long periods. The best candidate for the spin period with 1σ uncertainty is found at 17.34529(55) s. Since this period has a significance of 3.3σ and could not be confirmed in the other observations, probably owing to the lower number of detected counts, the spin period needs further confirmation.

3. Analyses and results of optical observations

An *I*-band finding chart from OGLE III is presented in Fig. 2. An optical counterpart, GSC 09142-02001, is found with an angular separation to the XMM-*Newton* position of $1.3''$ for the entry in the photometric catalogue of Massey (2002), giving $U = 14.48$ mag, $B = 15.39$ mag, $V = 15.45$ mag, and $R = 15.42$ mag as measured on MJD 51186. The NIR counterpart, 2MASS 01232743-7321223 (Skrutskie et al. 2006), is found with an angular separation of $1.1''$ with $J = 15.32$ mag, $H = 14.82$ mag, and $K_S = 15.06$ mag measured on MJD 51034. The corresponding OGLE sources are OGLE III SMC121.250




Fig. 3. Optical and NIR photometry of RX J0123.4-7321 dereddened and compared with normalised Kurucz model atmospheres for a B1 III and B3 III star. Labels give filters.




Fig. 5. Lomb-Scargle periodogram of the *I*-band light curve of RX J0123.4-7321 from OGLE III data before MJD 55000.




Fig. 4. Optical *I*-band light curve of RX J0123.4-7321 from OGLE III (before MJD 55000) and OGLE IV (after MJD 55000). Predicted outburst maxima are indicated by the dotted blue lines. The time of the XMM-Newton observation is marked by the dashed red line.




Fig. 6. OGLE III light curve of RX J0123.4-7321 for MJD < 55000 folded with the 119.87 d period. Phase = 0.5 corresponds to MJD 54387.14. The phases of the dates of the X-ray observations are indicated by dashed lines (see Table 1).

and OGLE IV SMC733.26.24 with a separation of $1.2''$ to the X-ray source. The OGLE *I*-band value of $I \approx 15.4$ mag is a typical magnitude of the variable light curve (see below). The magnitudes and colours are characteristic of a B star in the SMC. From optical spectroscopy, Evans et al. (2004) classified the source (2dFS 3624) as B1-3 III star. Since the density of early-type stars is relatively low in the wing of the SMC, a correlation with the X-ray source by chance is unlikely.

We dereddened these magnitudes to correct for a Galactic foreground of $E_{B-V} = 0.07$ mag (Schwering & Israel 1991) and compare them to a Kurucz model atmosphere (Kurucz 1979; Howarth 2011) of a B1 III and B3 III star ($T_{\text{eff}} = 25\,000, 18\,000$ K and $\log(g) = 3.0$) in Fig. 3. The models have been normalised to the *B*-band measurement. To account for possible variability of the source and the non-simultaneous observations, we included an uncertainty of 0.2 mag for the *I* and NIR magnitudes. There is evidence for an NIR excess, likely caused by the circum-stellar decretion disc and supporting the Be classification of the star.




Fig. 7. OGLE III light curves of RX J0123.4-7321 for six individual outbursts. Predicted outburst maxima are indicated by the dotted blue lines.

The long-term *I*-band light curve from OGLE phase III and IV is presented in Fig. 4. Besides a long-term variability, regular outbursts are seen. The light curve is typical for Be stars. A Lomb-Scargle periodogram (Lomb 1976; Scargle 1982) of the OGLE III data confirms the variability with a period of 119.87 days (Fig. 5). We derive an ephemeris for the *I*-band outburst maximum of

$$\text{MJD} = 54387.14 \pm 0.10 + (119.9 \pm 2.5) \times N,$$

with N an integer. The folded *I*-band light curve is presented in Fig. 6. The average outburst duration (FWHM) is 30 days as derived from the folded light curve. However, the evolution can be different for individual outbursts. For several outbursts we find a double-peaked shape, as also indicated in the folded light curve. Other outbursts only show one peak, where the maximum can be ~ 15 days before or after the expected time of the outburst maximum. Examples for individual outburst are presented in Fig. 7. In other BeXRBs, typically a fast rise and exponential decay is observed during optical outbursts (Bird et al. 2012). In general, individual outburst profiles will depend on NS-orbit parameters and perturbations of the decretion disc (Okazaki 1991).

All X-ray observations were done shortly after the times of possible optical outbursts, as indicated by the vertical lines in Fig. 6. If both optical and X-ray outburst appear during periastron passage, we would expect quasi-simultaneous outbursts as e.g. observed in AX J0058-720 (Haberl & Pietsch 2007) and IGR J05414-6858 (Sturm et al. 2012b). However, we note that the X-ray observations have been performed during times without OGLE coverage where the occurrence of optical outbursts is uncertain. The recently observed decrease in the *I*-band light curve and the vanishing of the outbursts can be caused by a loss of the decretion disc around the Be star. Interestingly, first the outbursts were missing after MJD 55400, although the quiescent *I*-band flux was at a similar level when outbursts have been observed around MJD 53000. Between MJD 55700 and 56000, a drop of the *I*-band flux was observed, followed by a brightening, probably due to a recovering of the decretion disc.

We note that RX J0123.4-7321 is along the line of sight to the H II region DEMS 160 (Davies et al. 1976), see Fig. 8. However, due to the high angular separation of $\sim 1'$ to the centre of DEMS 160, which corresponds to a distance of at least 17.5 pc, both sources are likely not connected directly to each other but might have formed during the same star-formation event.

4. Discussion and conclusions

We analysed ROSAT and OGLE data of RX J0123.4-7321. Due to the high X-ray variability by a factor of at least ~ 150 and the properties of the optical counterpart, we identify RX J0123.4-7321 as new BeXRB in the wing of the SMC. The *I*-band light curve of the optical counterpart shows a typical behaviour for a Be star in a BeXRB. The periodic outbursts are likely caused by binarity and reveal the orbit period of the NS of $P_{\text{orb}} = 119.78$ d. This places RX J0123.4-7321 in the sample of long-orbit systems (Knigge et al. 2011). From the Corbet relation (Corbet 1984; Corbet et al. 2009), we can roughly estimate the spin period of the neutron star to be between 10 and 800 s. Therefore, the source is unlikely to be identified with the 2.16 s pulsar XTE J0119-731 (Corbet et al. 2003) that was found by RXTE with an angular separation of $18'$. No indication of this period was found in the X-ray periodograms of RX J0123.4-7321. Further X-ray observations are necessary of the system in outburst, to confirm the possible spin period of 17.3 s and the




Fig. 8. Continuum-subtracted emission-line image of DEMS 160 from MCELS (Winkler et al. 2005). Red/green/blue give logarithmically scaled intensities for $H\alpha$ /[S II]/[O III]. RX J0123.4-7321 is marked with a white cross.

neutron-star nature of the compact object. If the current low-state of the *I*-band emission is caused by a loss of the decretion disc, no X-ray activity is expected. Follow-up X-ray observations are most likely successful, during new *I*-band outbursts, which indicate interaction of the NS and the decretion disc. Optical outbursts might reappear with brightening *I*-band emission and will be found in future OGLE monitoring. The system is included in the X-Ray variables OGLE Monitoring (XROM, Udalski 2008) system.

Acknowledgements. The OGLE project has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no. 246678 to AU. RS acknowledges support from the BMWI/DLR grant FKZ 50 OR 0907.

References

- Antoniu, V., Zezas, A., Hatzidimitriou, D., & Kalogera, V. 2010, *ApJ*, 716, L140
- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17
- Bird, A. J., Coe, M. J., McBride, V. A., & Udalski, A. 2012, *MNRAS*, 423, 3663
- Blackburn, J. K. 1995, in *Astronomical Society of the Pacific Conference Series*, Vol. 77, *Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 367
- Buchner, R., Bennett, K., Bignami, G. F., et al. 1983, *A&A*, 128, 245
- Chakrabarty, D., Levine, A. M., Clark, G. W., & Takeshima, T. 1998, *IAU Circ.*, 7048, 1
- Clark, G. W., Remillard, R. A., & Woo, J. W. 1997, *ApJ*, 474, L111
- Coe, M. J., Edge, W. R. T., Galache, J. L., & McBride, V. A. 2005, *MNRAS*, 356, 502
- Coe, M. J., Haberl, F., Sturm, R., et al. 2012, *MNRAS*, 424, 282
- Coe, M. J., Haberl, F., Sturm, R., et al. 2011, *MNRAS*, 414, 3281
- Corbet, R., Markwardt, C. B., Marshall, F. E., et al. 2003, *IAU Circ.*, 8064, 4
- Corbet, R. H. D. 1984, *A&A*, 141, 91
- Corbet, R. H. D., Coe, M. J., McGowan, K. E., et al. 2009, in *IAU Symposium*, Vol. 256, *IAU Symposium*, ed. J. T. van Loon & J. M. Oliveira, 361–366
- Davies, R. D., Elliott, K. H., & Meaburn, J. 1976, *MmRAS*, 81, 89
- Evans, C. J., Howarth, I. D., Irwin, M. J., Burnley, A. W., & Harries, T. J. 2004, *MNRAS*, 353, 601
- Gregory, P. C. & Lored, T. J. 1996, *ApJ*, 473, 1059
- Haberl, F., Eger, P., & Pietsch, W. 2008, *A&A*, 489, 327
- Haberl, F., Filipović, M. D., Pietsch, W., & Kahabka, P. 2000, *A&AS*, 142, 41
- Haberl, F. & Pietsch, W. 2004, *A&A*, 414, 667
- Haberl, F. & Pietsch, W. 2007, *A&A*, 476, 317
- Haberl, F., Sturm, R., Ballet, J., et al. 2012a, *A&A*, 545, A128

- Haberl, F., Sturm, R., Filipović, M. D., Pietsch, W., & Crawford, E. J. 2012b, *A&A*, 537, L1
- Haberl, F. & Zavlin, V. E. 2002, *A&A*, 391, 571
- Harris, J. & Zaritsky, D. 2004, *AJ*, 127, 1531
- Hénault-Brunet, V., Oskinova, L. M., Guerrero, M. A., et al. 2012, *MNRAS*, 420, L13
- Howarth, I. D. 2011, *MNRAS*, 413, 1515
- Knigge, C., Coe, M. J., & Podsiadlowski, P. 2011, *Nature*, 479, 372
- Kurucz, R. L. 1979, *ApJS*, 40, 1
- Laycock, S., Zezas, A., Hong, J., Drake, J. J., & Antoniou, V. 2010, *ApJ*, 716, 1217
- Li, K. L., Kong, A. K. H., Charles, P. A., et al. 2012, *ApJ*, 761, 99
- Liu, Q. Z., van Paradijs, J., & van den Heuvel, E. P. J. 2005, *A&A*, 442, 1135
- Lomb, N. R. 1976, *Ap&SS*, 39, 447
- Massey, P. 2002, *ApJS*, 141, 81
- Muller, E., Staveley-Smith, L., Zealey, W., & Stanimirović, S. 2003, *MNRAS*, 339, 105
- Okazaki, A. T. 1991, *PASJ*, 43, 75
- Okazaki, A. T. 2001, *PASJ*, 53, 119
- Reig, P. 2011, *Ap&SS*, 332, 1
- Scargle, J. D. 1982, *ApJ*, 263, 835
- Schwering, P. B. W. & Israel, F. P. 1991, *A&A*, 246, 231
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Stanimirovic, S., Staveley-Smith, L., Dickey, J. M., Sault, R. J., & Snowden, S. L. 1999, *MNRAS*, 302, 417
- Sturm, R., Haberl, F., Coe, M. J., et al. 2011, *A&A*, 527, A131
- Sturm, R., Haberl, F., Pietsch, W., et al. 2013, *A&A*, submitted
- Sturm, R., Haberl, F., Pietsch, W., et al. 2012a, *A&A*, 537, A76
- Sturm, R., Haberl, F., Rau, A., et al. 2012b, *A&A*, 542, A109
- Townsend, L. J., Coe, M. J., Corbet, R. H. D., et al. 2011, *MNRAS*, 410, 1813
- Tozzi, P., Gilli, R., Mainieri, V., et al. 2006, *A&A*, 451, 457
- Trümper, J. 1982, *Advances in Space Research*, 2, 241
- Udalski, A. 2008, *Acta Astron.*, 58, 187
- Udalski, A., Szymanski, M. K., Soszynski, I., & Poleski, R. 2008, *Acta Astron.*, 58, 69
- Winkler, P. F., Young, A. L., Braziunas, D., et al. 2005, in *Bulletin of the American Astronomical Society*, Vol. 37, American Astronomical Society Meeting Abstracts, 132.03
- Ziolkowski, J. 2002, *Mem. Soc. Astron. Italiana*, 73, 1038